

Minimum soapy union

Francois Nicolas¹ and Sebastian Böcker¹

Lehrstuhl für Bioinformatik, Friedrich-Schiller-Universität Jena, Ernst-Abbe-Platz 2,
Jena, Germany, {francois.nicolas, sebastian.boecker}@uni-jena.de

Abstract. Define MINIMUM SOAPY UNION (MINSU) as the following optimization problem: given a k -tuple $(X_1, X_2, \dots, X_k)$ of finite integer sets, find a k -tuple $(t_1, t_2, \dots, t_k)$ of integers that minimizes the cardinality of $(X_1 + t_1) \cup (X_2 + t_2) \cup \dots \cup (X_k + t_k)$. We show that MINSU is NP-complete, APX-hard, and polynomial for fixed k .

MINSU appears naturally in the context of protein shotgun sequencing: Here, the protein is cleaved into short and overlapping peptides, which are then analyzed by tandem mass spectrometry. To improve the quality of such spectra, one then asks for the mass of the unknown prefix (the shift) of the spectrum, such that the resulting shifted spectra show a maximum agreement. For real-world data the problem is even more complicated than our definition of MINSU; but our intractability results clearly indicate that it is unlikely to find a polynomial time algorithm for shotgun protein sequencing.

1 Introduction

The aim of this paper is to study the computational complexity of the following optimization problem:

Name: MINIMUM SOAPY UNION (MINSU)

Input: a finite set A and an indexed family $(X_a)_{a \in A}$ of non-empty finite sets of rational integers.

Solution: an indexed family $(t_a)_{a \in A}$ of rational integers.

Measure: the cardinality of $\bigcup_{a \in A} (X_a + t_a)$.

Let us name SOAPY UNION (SU) the decision problem associated with MINSU. The names have been chosen by analogy with the SOAPY SET COVER problem [14]. Clearly, SU is a *number problem* [11]. MINSU can be seen as a generalization of the SUBSET MATCHING problem [8]: optimally solving SUBSET MATCHING is equivalent to optimally solving the restriction of MINSU to those instances $(X_a)_{a \in A}$ such that the cardinality of A equals 2.

MINSU naturally appears in the context of protein shotgun sequencing [6,5,4]. (This problem must not be confused with the more widely known peptide shotgun sequencing.) Sequencing the protein means that we want to determine its amino acid sequence. We assume that no genomic information is available for the protein, so that its sequence cannot be derived from the genomic

information. This is the case for many proteins even in humans, monoclonal antibodies being an important example [5]. Experimentally, the protein is cleaved into short and overlapping peptides, which are then analyzed by tandem mass spectrometry. To improve the quality of such spectra, one then asks for the mass of the unknown prefix (the shift) of the spectrum, such that the resulting shifted spectra show a maximum agreement. For real-world data the problem is even more complicated than our definition of MINSU; but our intractability results clearly indicate that it is unlikely to find a polynomial time algorithm for shotgun protein sequencing.

Contribution. In Section 2, we prove that SU belongs to NP and that MINSU can be solved in polynomial time for fixed A . In Section 3, we show that SU is strongly NP-hard; furthermore, we prove that there exists a real number $\rho > 1$ such that if MINSU is ρ -approximable in pseudo-polynomial time then $P = NP$.

Notation and definitions. For every finite set S , $|S|$ denotes the cardinality of S . For all sets A and S , S^A denotes the set of all families of elements of S indexed by A .

The ring of rational integers is denoted $\mathbb{Z}$. For every integer $n \geq 0$, $[1, n]$ denotes the set of all $k \in \mathbb{Z}$ such that $1 \leq k \leq n$.

A(n undirected) *graph* is a pair $G = (V, E)$, where V is a finite set and E is a set of 2-element subsets of V : the elements of V are the *vertices* of G , the elements of E are the *edges* of G , and for each edge $e \in E$, the elements of e are the *extremities* of e .

Let MIN be a minimization problem. The *decision problem associated with* MIN is: given an instance I of MIN and an integer $k \geq 0$, decide whether there exists a solution of MIN on I with measure at most k .

2 Membership

For each instance $(X_a)_{a \in A}$ of MINSU, the set of all feasible solutions of MINSU on $(X_a)_{a \in A}$ equals $\mathbb{Z}^A$, which is infinite. Therefore, MINSU is not an NP-*optimization problem* [2], and thus the membership of SU in NP is not completely trivial.

Let $G = (V, E)$ be a graph. A *disconnection* of G is a pair (B, C) such that $B \neq \emptyset$, $C \neq \emptyset$, $B \cap C = \emptyset$, $V = B \cup C$, and for every $(b, c) \in B \times C$, $\{b, c\} \notin E$. A graph is called *disconnected* if it admits a disconnection. A graph that is not disconnected is called *connected*.

Let $(Y_a)_{a \in A}$ be an indexed family of sets. The *intersection graph* of $(Y_a)_{a \in A}$ is defined as follows: its vertex set equals A and for all $b, c \in A$ with $b \neq c$, $\{b, c\}$ is one of its edges if, and only if, $Y_b \cap Y_c \neq \emptyset$.

Lemma 1. *Let $(Y_a)_{a \in A}$ be an instance of MINSU. If the intersection graph of $(Y_a)_{a \in A}$ is disconnected then there exists $(u_a)_{a \in A} \in \mathbb{Z}^A$ such that*

$$\left| \bigcup_{a \in A} (Y_a + u_a) \right| < \left| \bigcup_{a \in A} Y_a \right|. \quad (1)$$

Proof. For each subset $B \subseteq A$, put $Y_B = \bigcup_{b \in B} Y_b$. Let (B, C) be a disconnection of the intersection graph of $(Y_a)_{a \in A}$. Let $r \in Y_B$ and $s \in Y_C$ be fixed. Set $u_b = -r$ for every $b \in B$ and $u_c = -s$ for every $c \in C$. On the one hand, we have $Y_B \cap Y_C = \emptyset$ and $Y_A = Y_B \cup Y_C$, so

$$|Y_A| = |Y_B| + |Y_C|. \quad (2)$$

On the other hand, we have

$$\bigcup_{a \in A} (Y_a + u_a) = (Y_B - r) \cup (Y_C - s)$$

and

$$(Y_B - r) \cap (Y_C - s) \neq \emptyset$$

because $0 \in (Y_B - r) \cap (Y_C - s)$; it follows

$$\left| \bigcup_{a \in A} (Y_a + u_a) \right| < |Y_B| + |Y_C|. \quad (3)$$

It now suffices to combine Equations (2) and (3) to obtain Equation (1). $\square$

Lemma 1 can be restated as follows:

Lemma 2. *Let $(X_a)_{a \in A}$ be an instance of MINSU. For any optimum solution $(t_a)_{a \in A}$ of MINSU on $(X_a)_{a \in A}$, the intersection graph of $(X_a + t_a)_{a \in A}$ is connected.*

Proof. Let $(t_a)_{a \in A} \in \mathbb{Z}^A$ be such that the intersection graph of $(X_a + t_a)_{a \in A}$ is disconnected. Set $Y_a = X_a + t_a$ for each $a \in A$. By Lemma 1, there exists $(u_a)_{a \in A} \in \mathbb{Z}^A$ such that Equation (1) holds. It follows that $(t_a + u_a)_{a \in A}$ is a better solution of MINSU on $(X_a)_{a \in A}$ than $(t_a)_{a \in A}$. $\square$

Definition 1. *Let $G = (V, E)$ be a graph. Put*

$$\tilde{E} = \{(a, b) \in V \times V : \{a, b\} \in E\}.$$

An antisymmetric edge-weight function on G is a function ϖ from $\tilde{E}$ to $\mathbb{Z}$ such that $\varpi(b, c) = -\varpi(c, b)$ for every $(b, c) \in \tilde{E}$. For each antisymmetric edge-weight function ϖ on G , define $S(G, \varpi)$ as the set of all $(t_a)_{a \in V} \in \mathbb{Z}^V$ such that $t_b - t_c = \varpi(b, c)$ for all $(b, c) \in \tilde{E}$.

Let us comment Definition 1. The function ϖ assigns both a magnitude and an orientation to each edge of G : for all $a, b \in V$ such that $\{a, b\} \in E$, the magnitude of $\{a, b\}$ is the absolute value of $\varpi(a, b)$ and the orientation of $\{a, b\}$ is determined by the sign of $\varpi(a, b)$. It is clear that for every $(t_a)_{a \in V} \in S(G, \varpi)$ and every $u \in \mathbb{Z}$, $(t_a + u)_{a \in V} \in S(G, \varpi)$. If G is connected then either $S(G, \varpi)$ is empty or there exists $(t_a)_{a \in V} \in \mathbb{Z}^V$ such that $S(G, \varpi) = \{(t_a + u)_{a \in V} : u \in \mathbb{Z}\}$. If G is connected and $S(G, \varpi) \neq \emptyset$ then for any $(b, u) \in V \times \mathbb{Z}$, the unique element

$(t_a)_{a \in V} \in S(G, \varpi)$ that satisfies $t_b = u$ is computable from G, ϖ, b , and u in polynomial time. A *closed walk* in G is a finite sequence $(a_0, a_1, a_2, \dots, a_k)$ such that $a_0 = a_k$ and $\{a_{i-1}, a_i\} \in E$ for every $i \in [1, k]$; the *weight* of $(a_0, a_1, a_2, \dots, a_k)$ under ϖ is defined as $\varpi(a_0, a_1) + \varpi(a_1, a_2) + \dots + \varpi(a_{k-1}, a_k)$. A (simple) *cycle* in G is a closed walk $(a_0, a_1, a_2, \dots, a_k)$ in G such that for all $i, j \in [1, k]$, $a_i = a_j$ implies $i = j$. The following three conditions are equivalent:

1. The set $S(G, \varpi)$ is non-empty.
2. The weight under ϖ of every closed walk in G equals 0.
3. The weight under ϖ of every cycle in G equals 0.

The second and third conditions can be thought as abstract forms of Kirchhoff's voltage law.

A *tree* is a connected graph with one fewer edges than vertices, or equivalently, an acyclic connected graph. An arbitrary graph $G = (V, E)$ is connected if, and only if, there exists a subset $E' \subseteq E$ such that (V, E') is a tree ((V, E') is then called a *spanning tree* of G).

Lemma 3. *Let $(X_a)_{a \in A}$ be an instance of MINSU. There exist a tree H with vertex set A and an antisymmetric edge-weight function ϖ on H that satisfy the following two conditions:*

1. *Every integer in the range of ϖ can be written as the difference of two elements of $\bigcup_{a \in A} X_a$.*
2. *Every element of $S(H, \varpi)$ is an optimum solution of MINSU on $(X_a)_{a \in A}$.*

Proof. Let $(t_a)_{a \in A}$ be an optimum solution of MINSU on $(X_a)_{a \in A}$. Let H be a spanning tree of the intersection graph of $(X_a + t_a)_{a \in A}$: such a tree exists by Lemma 2. Let ϖ be the antisymmetric edge-weight function on H defined by: for all $b, c \in A$ such that $\{b, c\}$ is an edge of H , $\varpi(b, c) = t_b - t_c$.

For all $b, c \in A$, such that $\{b, c\}$ is an edge of the intersection graph of $(X_a + t_a)_{a \in A}$, $(X_b + t_b) \cap (X_c + t_c)$ is non-empty, and thus $t_b - t_c$ belongs to $X_c - X_b$. Therefore, the first condition holds. Now, remark that $S(H, \varpi) = \{(t_a + u)_{a \in A} : u \in \mathbb{Z}\}$, so the second condition holds. $\square$

Theorem 1. *SU belongs to NP.*

Proof. Let $((X_a)_{a \in A}, k)$ be an arbitrary instance of SU. We propose the following (non-deterministic) algorithm to decide whether $((X_a)_{a \in A}, k)$ is a yes-instance of SU:

- Guess a tree H with vertex set A and an antisymmetric edge-weight function ϖ on H such that the first condition of Lemma 3 holds.
- Compute an element $(t_a)_{a \in A} \in S(H, \varpi)$.
- Check whether the cardinality of $\bigcup_{a \in A} (X_a + t_a)$ is at most k .

By Lemma 3, the algorithm is correct. Moreover, the bit-length of the guess (*i.e.*, the ordered pair (H, ϖ)) is polynomial in the bit-length of the input (*i.e.*, the instance $(X_a)_{a \in A}$), so the algorithm can be implemented in non-deterministic polynomial time. $\square$

Let m be a positive integer and let X be a subset of $\mathbb{Z}$ such that $X = -X$. On each given m -edge graph, there are exactly $|X|^m$ distinct antisymmetric edge-weight functions whose ranges are subsets of X .

Let n be a positive integer and let $\mathcal{T}_n$ denote the set of all trees with vertex set $[1, n]$. *Cayley's formula* ensures $|\mathcal{T}_n| = n^{n-2}$ [12]. Moreover, every tree can be reconstructed in polynomial time from its *Prüfer code* [12], so $\mathcal{T}_n$ is enumerable in $O(n^{O(n)})$ time.

Theorem 2. *There exists an algorithm that, for each instance $(X_a)_{a \in A}$ of MINSU given as input, returns an optimum solution of MINSU on $(X_a)_{a \in A}$ in $O(N^{O(|A|)})$ time, where N denotes the bit-length of $(X_a)_{a \in A}$.*

Proof. Put $U = \bigcup_{a \in A} X_a$. Let $\mathcal{H}$ denote the set of all ordered pairs of the form (H, ϖ) , where H is a tree with vertex set A and ϖ is an antisymmetric edge-weight function on H whose range is a subset of $U - U$. We propose the following algorithm to solve MINSU on $(X_a)_{a \in A}$:

- For each $(H, \varpi) \in \mathcal{H}$, compute an element of $S(H, \varpi)$.
- Return a best solution of MINSU on $(X_a)_{a \in A}$ among those computed at the previous step.

By Lemma 3, the algorithm returns an optimum solution of MINSU on $(X_a)_{a \in A}$. Moreover, remark that $|\mathcal{H}| = |A|^{|A|-2} |U - U|^{|A|-1}$ and that $\mathcal{H}$ is enumerable in $O(N^{O(|A|)})$ time. Therefore, the algorithm can be implemented to run in $O(N^{O(|A|)})$ time. $\square$

3 Hardness

The aim of this section is prove the hardness results for MINSU.

Let $G = (V, E)$ be a graph. A *vertex cover* of G is a subset $C \subseteq V$ such that $C \cap e \neq \emptyset$ for every $e \in E$: a vertex cover is a subset of vertices that contains at least one extremity of each edge.

Name: MINIMUM VERTEX COVER (MINVC)

Input: a graph G .

Solution: a vertex cover C of G .

Measure: the cardinality of C .

The decision problem associated with MINVC is named VERTEX COVER (VC). It is well-known that VC is NP-complete [11].

To prove that SU is (strongly) NP-complete, we show that VC Karp-reduces to (a suitable restriction of) SU. The following gadget plays a crucial role in our reduction as well as in other reductions that can be found in the literature [14,13]:

Definition 2. *For each integer $n \geq 1$, define $R_n = \{(i-1)n^2 + i^2 : i \in [1, n]\}$.*

A *Golomb ruler* [10,15,3] is a finite subset $R \subseteq \mathbb{Z}$ that satisfies the following three equivalent conditions:

- For every $t \in \mathbb{Z}$, $t \neq 0$ implies $|R \cap (R + t)| \leq 1$.
- For every integer $d > 0$, there exists at most one $(r, s) \in R \times R$ such that $r - s = d$.
- For all $r_1, r_2, s_1, s_2 \in R$, $r_1 + r_2 = s_1 + s_2$ implies $\{r_1, r_2\} = \{s_1, s_2\}$.

Actually, only the first condition is referred to in what follows. Among other convenient properties our gadget sets are Golomb rulers:

Lemma 4. *Let n be a positive integer. The following four properties hold.*

1. *The least element of R_n is 1 and the greatest element of R_n is n^3 .*
2. *The cardinality of R_n equals n .*
3. *The distance between any two elements of R_n is at least $n^2 + 3$.*
4. *R_n is a Golomb ruler.*

Proof. Properties 1 and 2 are clear. Proofs of Property 4 can be found in [14,13]. Finally, remark that for every $i \geq 1$, we have

$$(in^2 + (i+1)^2) - ((i-1)n^2 + i^2) = n^2 + 2i + 1 \geq n^2 + 3.$$

Hence, the distance between any two consecutive elements of R_n is at least $n^2 + 3$, and thus Property 3 holds. $\square$

Theorem 3. *SU is strongly NP-hard.*

Proof. Put $f(x) = (\frac{1}{4}x + 2)^3 + \frac{1}{2}x - 4$. Let AUX denote the restriction of SU to those instances $((X_a)_{a \in A}, k)$ such that the absolute value of every integer in $\{k\} \cup \bigcup_{a \in A} X_a$ is at most $f(\max_{a \in A} |X_a|)$. We prove that AUX is NP-hard which implies the theorem. More precisely, we show that VC Karp-reduces to AUX.

Presentation of the reduction. Let I be an arbitrary instance of VC. The reduction maps I to an instance J of SU that is defined as follows. Let G, V, E , and k be such that $I = (G, k)$ and $G = (V, E)$. Let n denote the cardinality of V . Without loss of generality, we may assume $V = [1, n]$ and $k < n$ because I is a yes-instance of VC whenever $k \geq n$. Let $(y_e)_{e \in E}, (z_e)_{e \in E} \in V^E$ be such that $e = \{y_e, z_e\}$ for every $e \in E$. Set

$$\begin{aligned} A &= \{\emptyset\} \cup E, \\ s &= (n+4)^3, \\ R &= R_{n+4}, \\ X_{\emptyset} &= (V - s - n) \cup (R - s) \cup (R + n) \cup (V + s + n), \\ X_e &= \{z_e - n\} \cup R \cup \{y_e + s\} \end{aligned}$$

for each $e \in E$, and

$$J = ((X_a)_{a \in A}, |X_{\emptyset}| + k).$$

Clearly, J is computable from I in polynomial time.

An instance of AUX. Let us prove that J is in fact an instance of AUX. With the help of Lemmas 4.1 and 4.2, it is easy to see that $1 - s - n$ is the least element of $\bigcup_{a \in A} X_a$, that $s + 2n$ is the greatest element of $\bigcup_{a \in A} X_a$, and that the cardinality of $X_\emptyset$ equals $4n + 8$. The latter property implies $|X_\emptyset| + k < 5n + 8$. Hence, the absolute value of every integer in $\{|X_\emptyset| + k\} \cup \bigcup_{a \in A} X_a$ is at most $s + 2n$. Now, remark that $s + 2n = f(|X_\emptyset|) \leq f(\max_{a \in A} |X_a|)$.

Correctness of the reduction. It remains to prove that I is a yes-instance of VC if, and only if, J is a yes-instance of SU.

Lemma 5. *For every $e \in E$, it holds true that*

1. $(X_e - s) \setminus X_\emptyset = \{y_e\}$ and that
2. $(X_e + n) \setminus X_\emptyset = \{z_e\}$.

Proof. We only prove Property 1 because Property 2 can be proven in the same way. Put $Y = \{z_e - s - n\} \cup (R - s)$. It is clear that $X_e - s = Y \cup \{y_e\}$ and $Y \subseteq X_\emptyset$. Therefore, we have

$$X_e - s = \{y_e\} \setminus X_\emptyset. \quad (4)$$

Moreover, it follows from Lemma 4.1 that

- the greatest element of $(V - s - n) \cup (R - s)$ equals 0 and that
- the least element of $(R + n) \cup (V + s + n)$ equals $n + 1$.

Therefore, $X_\emptyset$ does not contain any element of $[1, n]$. In particular, y_e does not belong to $X_\emptyset$. Combining the latter fact with Equation (4), we obtain Property 1. $\square$

Lemma 6. *For every $t \in \mathbb{Z}$, $|(R + t) \setminus X_\emptyset| < n$ implies $t \in \{-s, +n\}$.*

Proof. Let us first bound from above the cardinality of $(R + t) \cap X_\emptyset$. For each $\tau \in \mathbb{Z}$, put $P_\tau = (R + t) \cap (V + \tau)$ and $Q_\tau = (R + t) \cap (R + \tau)$. First, it follows from Lemma 4.3 that $|P_\tau| \leq 1$. Second, $\tau \neq t$ implies $|Q_\tau| \leq 1$ by Lemma 4.4. And third, it holds that

$$(R + t) \cap X_\emptyset = P_{-s-n} \cup Q_{-s} \cup Q_n \cup P_{s+n}.$$

Now, assume $t \notin \{-s, +n\}$. From the preceding three facts, we deduce that

$$|(R + t) \cap X_\emptyset| \leq |P_{-s-n}| + |Q_{-s}| + |Q_n| + |P_{s+n}| \leq 4.$$

(In fact, it is not hard to see that $|(R + t) \cap X_\emptyset| \leq 2$ holds: $t > +n$ implies $P_{-s-n} = Q_{-s} = \emptyset$, $-s < t < +n$ implies $P_{-s-n} = P_{s+n} = \emptyset$, and $t < -s$ implies $Q_n = P_{s+n} = \emptyset$.) Since $|R + t| = n + 4$ by Lemma 4.2, we finally get

$$|(R + t) \setminus X_\emptyset| = n + 4 - |(R + t) \cap X_\emptyset| \geq n.$$

$\square$

(If). Assume that I is a yes-instance of VC. Then, there exists a vertex cover C of G with $|C| \leq k$. Put $F = \{e \in E : y_e \in C\}$. Set $t_\emptyset = 0$, $t_e = -s$ for each $e \in F$, and $t_e = +n$ for each $e \in E \setminus F$. On the one hand, it holds that

$$\left| \bigcup_{a \in A} (X_a + t_a) \right| = |X_\emptyset| + \left| \bigcup_{e \in E} (X_e + t_e) \setminus X_\emptyset \right| \quad (5)$$

because $t_\emptyset = 0$. On the other hand, it follows from Lemma 5 that

$$\bigcup_{e \in E} (X_e + t_e) \setminus X_\emptyset = \{y_e : e \in F\} \cup \{z_e : e \in E \setminus F\}. \quad (6)$$

Since the right-hand side of Equation (6) is a subset of C , we have

$$\left| \bigcup_{e \in E} (X_e + t_e) \setminus X_\emptyset \right| \leq k. \quad (7)$$

We then get

$$\left| \bigcup_{a \in A} (X_a + t_a) \right| \leq |X_\emptyset| + k \quad (8)$$

by combining Equations (5) and (7). Hence, J is a yes-instance of SU.

(Only if). Assume that J is a yes-instance of SU. Then, there exists $(t_a)_{a \in A} \in \mathbb{Z}^A$ such that Equation (8) holds. Replacing $(t_a)_{a \in A}$ with $(t_a - t_\emptyset)_{a \in A}$ leaves the cardinality of $\bigcup_{a \in A} (X_a + t_a)$ unchanged; therefore, we may assume that $t_\emptyset = 0$; in particular, Equation (5) holds.

Put

$$C = \bigcup_{e \in E} (X_e + t_e) \setminus X_\emptyset.$$

Combining Equations (5) and (8), we obtain Equation (7), or equivalently, $|C| \leq k$. Now, let us prove that C is a vertex cover of G . Consider an arbitrary edge $e \in E$. Since we have

$$(R + t_e) \setminus X_\emptyset \subseteq (X_e + t_e) \setminus X_\emptyset \subseteq C,$$

it follows from Lemma 6 that $t_e \in \{-s, +n\}$. Consequently, Lemma 5 ensures that some extremity of e belongs to $(X_e + t_e) \setminus X_\emptyset$, and this extremity is *a fortiori* in C . Hence, I is a yes-instance of VC. $\square$

A graph $G = (V, E)$ is called *cubic* if for every vertex $v \in V$, the degree of v in G (i.e., the cardinality of $\{w \in V : \{v, w\} \in E\}$) equals 3. Let MINVC3 denote the restriction of MINVC to cubic graphs. MINVC3 is APX-complete under L-reduction [1]; moreover, if MINVC3 is $\frac{100}{99}$ -approximable in polynomial time then $P = NP$ [7].

To prove that MINSU is “strongly” APX-hard, which is a better result than Theorem 3, we show that MINVC3 L-reduces to a suitable restriction of MINSU. In fact, we simply adapt the proof of Theorem 3.

Theorem 4. *There exists a real constant $\rho > 1$ such that if MINSU is ρ -approximable in pseudo-polynomial time then $P = NP$.*

Proof. Let f be as in the proof of Theorem 3 and let MINAUX denote the restriction of MINSU to those instances $(X_a)_{a \in A}$ such that the absolute value of every integer in $\bigcup_{a \in A} X_a$ is at most $f(\max_{a \in A} |X_a|)$. We prove that MINAUX is APX-hard, which implies the theorem because every pseudo-polynomial-time approximation algorithm for MINSU is a polynomial-time approximation algorithm for MINAUX. More precisely, we show that MINVC3 L-reduces [2] to MINAUX. We use the notation of the proof of Theorem 3.

From a graph to an instance of AUX. Let τ denote the minimum cardinality of a vertex cover of G . Let v denote the minimum cardinality of $\bigcup_{a \in A} (X_a + t_a)$ over all $(t_a)_{a \in A} \in \mathbb{Z}^A$. Clearly, $(X_a)_{a \in A}$ is computable from G in polynomial time ($(X_a)_{a \in A}$ is independent of k), $(X_a)_{a \in A}$ is an instance of MINAUX, and $v = |X_\emptyset| + \tau = 4n + 8 + \tau$.

Now, assume that G is cubic and $n \geq 24$. The first assumption implies $3\tau \geq |E| \geq n$. It follows

$$4n + 8 = \left(4 + \frac{8}{n}\right)n \leq \left(12 + \frac{24}{n}\right)\tau \leq 13\tau,$$

and thus $v \leq 14\tau$.

From a solution of AUX to a vertex cover. Let $(t_a)_{a \in A} \in \mathbb{Z}^A$. Put

$$k = \left| \bigcup_{a \in A} (X_a + t_a) \right| - |X_\emptyset|.$$

There exists a vertex cover C of G that satisfies $|C| \leq k$, or equivalently,

$$|C| - \tau \leq \left| \bigcup_{a \in A} (X_a + t_a) \right| - v.$$

Moreover, such a vertex cover is computable from G and $(t_a)_{a \in A}$ in polynomial time:

- if $k \geq n$ then set $C = V$ and
- if $k < n$ then set $C = \bigcup_{e \in E} (X_e + t_e - t_\emptyset) \setminus X_\emptyset$.

Conclusion. Let ε be a positive real number. If MINSU is $(1 + \varepsilon)$ -approximable in pseudo-polynomial time then MINVC3 is $(1 + 14\varepsilon)$ -approximable in polynomial time. Therefore, if MINSU is $\frac{1387}{1386}$ -approximable in pseudo-polynomial time then $P = NP$. $\square$

An immediate corollary of Theorem 4 is that MINSU does not admit any (pseudo-)polynomial time approximation scheme.

4 Open questions

The following three questions remain open: Does there exist a constant $\rho > 1$ such that MINSU is ρ -approximable in (pseudo-)polynomial time? Is SU *fixed-parameter tractable* [9] with respect to parameter $|A|$? Is SU solvable in polynomial time for bounded $\max_{a \in A} |X_a|$?

References

1. P. Alimonti and V. Kann. Some APX-completeness results for cubic graphs. *Theor. Comput. Sci.*, 237(1-2):123–134, 2000.
2. G. Ausiello, P. Crescenzi, G. Gambosi, V. Kann, A. Marchetti-Spaccamela, and M. Protasi. *Complexity and Approximation: Combinatorial Optimization Problems and Their Approximability Properties*. Springer-Verlag, second edition, 2003.
3. W. C. Babcock. Intermodulation interference in radio systems: Frequency of occurrence and control by channel selection. *Bell Syst. Tech. J.*, 32(1):63–73, 1953.
4. N. Bandeira, J. Ng, D. Meluzzi, R. G. Linington, P. Dorrestein, and P. A. Pevzner. De novo sequencing of nonribosomal peptides. In *Proc. of Research in Computational Molecular Biology (RECOMB 2008)*, volume 4955 of *Lect. Notes Bioinform.*, pages 181–195. Springer, Berlin, 2008.
5. N. Bandeira, V. Pham, P. Pevzner, D. Arnott, and J. R. Lill. Automated de novo protein sequencing of monoclonal antibodies. *Nat. Biotechnol.*, 26(12):1336–1338, 2008.
6. N. Bandeira, H. Tang, V. Bafna, and P. Pevzner. Shotgun protein sequencing by tandem mass spectra assembly. *Anal. Chem.*, 76(24):7221–7233, 2004.
7. M. Chlebík and J. Chlebíková. Complexity of approximating bounded variants of optimization problems. *Theor. Comput. Sci.*, 354(3):320–338, 2006.
8. R. Clifford and B. Sach. Fast approximate point set matching for information retrieval. In *Proc. of Conference on Current Trends in Theory and Practice of Computer Science (SOFSEM 2007)*, volume 4362 of *Lect. Notes Comput. Sci.*, pages 212–223. Springer, Berlin, 2007.
9. J. Flum and M. Grohe. *Parameterized Complexity Theory*. Springer-Verlag, 2006.
10. M. Gardner. *Wheels, Life and Other Mathematical Amusements*. W. H. Freeman and Co., 1983.
11. M. R. Garey and D. S. Johnson. *Computers and Intractability: A Guide to the Theory of NP-Completeness*. W. H. Freeman and Co., 1979.
12. F. Harary and E. M. Palmer. *Graphical Enumeration*. Academic Press, 1973.
13. M. Michael, F. Nicolas, and E. Ukkonen. On the complexity of finding gapped motifs. *J. Discrete Algorithms*, 8(2):131–142, 2010.
14. F. Nicolas and É. Rivals. Hardness of optimal spaced seed design. *J. Comput. System Sci.*, 74(5):831–849, 2008.
15. S. Sidon. Ein Satz über trigonometrische Polynome und seine Anwendung in der Theorie der Fourier-Reihen. *Math. Ann.*, 106(1):536–539, 1932.